

0=Never (0%)	1=Rarely (20%)	2=Sometimes (40%)	3=Often (60%)	4=Very Often (80%)	5=Always (100%)
--------------	----------------	-------------------	---------------	--------------------	-----------------

Scores based on a self report scale for Children- presents the reduction in symptoms related to hyperactivity

Patient	Hyperactivity			Impulsive			Can't sit still		
	Before	After	Improve	Before	After	Improve	Before	After	Improve
6Y Male - ADD	40	20	50%	20	20	0%	40	20	50%
7Y Male - ADD	40	20	50%	20	0	100%	20	0	100%
7Y Male - ADHD	60	20	67%	60	20	67%	60	40	33%
8Y Female - ADD/LD	40	20	50%	0	0	0%	20	0	100%
8Y Male - ADHD	80	40	50%	100	40	60%	80	20	75%
10Y Male - ADD/ADP	60	20	67%	20	0	100%	20	20	0%
10Y Female - ADHD	100	40	60%	80	40	50%	80	20	75%
11Y Male - ADHD	80	40	50%	80	20	75%	80	20	75%
12Y Female - ADD	40	0	100%	20	0	100%	0	0	0%
13Y Female - ADD/LD	40	20	50%	0	0	0%	20	20	0%
14Y Male - ADD/ADP	40	20	50%	20	0	100%	20	0	100%
16Y Female - ADHD	60	20	67%	40	0	100%	40	0	100%
16Y Female - ADHD	80	40	50%	80	40	50%	80	20	75%
17Y Male - ADHD	100	40	60%	80	40	50%	80	40	50%

Patient	Average %	Average %
	Hyperactivity Before	Hyperactivity After
6Y Male - ADD	33	20
7Y Male - ADD	27	7
7Y Male - ADHD	60	27
8Y Female - ADD/LD	20	7
8Y Male - ADHD	87	33
10Y Male - ADD/ADP	33	13
10Y Female - ADHD	87	33
11Y Male - ADHD	80	27
12Y Female - ADD	20	0
13Y Female - ADD/LD	20	13
14Y Male - ADD/ADP	27	7
16Y Female - ADHD	47	7
16Y Female - ADHD	80	33
17Y Male - ADHD	87	40

Average entire group                      50                      19

Table 1    Presents reduction in hyperactivity symptoms

